

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 December 2000 (07.12.2000)

PCT

(10) International Publication Number
WO 00/72847 A1

- (51) International Patent Classification⁷: A61K 31/496, 9/10 (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).
- (21) International Application Number: PCT/IB00/00593 (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 8 May 2000 (08.05.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/136,268 27 May 1999 (27.05.1999) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): PFIZER PRODUCTS INC. [US/US]; Eastern Point Road, Groton, CT 06340 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): ARENSON, Daniel, Ray [US/US]; 16 Colony Road, East Lyme, CT 06333 (US). QI, Hong [US/US]; 4529 Concord Lane, Northbrook, IL 60062 (US).
- Published:
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: ZIPRASIDONE SUSPENSION

(57) Abstract: Compositions comprising ziprasidone free base or a difficult to wet pharmaceutically acceptable ziprasidone acid addition salt, a polysorbate, and colloidal silicon dioxide form good aqueous suspensions having a useful shelf life and are easily re-suspended if settling occurs.

WO 00/72847 A1


ZIPRASIDONE SUSPENSION

Field Of The Invention

This invention relates to an oral suspension comprising ziprasidone free base or a pharmaceutically acceptable ziprasidone acid addition salt, a polysorbate, colloidal silicon dioxide, a viscosity agent, and water. In a more specific aspect, the invention relates to such a suspension which is taste masked. The invention further relates to a method of treating a psychosis with such a formulation.

Background Of The Invention

Ziprasidone is a known compound having the structure:



It is disclosed in U.S. patents 4,831,031 and 5,312,925, both of which are herein incorporated by reference in their entirety, has utility as a neuroleptic, and is thus useful, *inter alia*, as an antipsychotic. It is typically administered orally as the hydrochloride acid addition salt, ziprasidone hydrochloride monohydrate. The hydrochloride salt is advantageous in that it is a high permeability drug, a factor which favorably affects bioavailability. The hydrochloride salt, as well as other ziprasidone acid addition salts, does, however, possess relatively poor aqueous solubility, a factor which unfavorably affects bioavailability.

Difficult to wet pharmaceutically acceptable compounds can be problematic in the pharmaceutical arts from a formulations perspective. For example, ziprasidone hydrochloride, in addition to having low solubility, is difficult to wet with an aqueous medium, and thereby presents special problems from the standpoint of trying to form an aqueous suspension. In the discussion which follows, ziprasidone hydrochloride is discussed as an exemplary member of the class composed of ziprasidone free base and ziprasidone acid addition salts which are difficult to wet. The invention is not to be taken as being limited to ziprasidone hydrochloride, however.

Owing to difficulties in wetting ziprasidone acid addition salts such as ziprasidone hydrochloride, the material is difficult to adequately suspend in an aqueous medium without having to resort to using long periods of high shear mixing. An ordinary laboratory homogenizer generally does not wet ziprasidone hydrochloride without being run for a very long time. Long blending periods almost inevitably lead to foaming and still yield poor results with drug aggregates still being visibly present, entrained in the foam. Thus, ziprasidone hydrochloride tends to float on the surface of water and other aqueous media and can be induced to form a suspension only with physical measures (high shear mixing for long times) that are considered extreme.

An alternative mixing procedure comprises first adding only a small amount of water to the ziprasidone salt, followed by grinding to wet the bulk drug substance. This wets the mass sufficiently so that it can be suspended in water. This procedure is still disadvantageous however in that it is difficult to scale up. Moreover, once ziprasidone hydrochloride has been induced to form an aqueous suspension in this fashion, an additional challenge is the prevention or retarding of rapid re-settling, which occurs relatively quickly with ziprasidone hydrochloride, generally within an hour or so, depending on the particle size. Settling can be particularly problematic for a pharmaceutical suspension since the suspension must be adequately re-suspended to ensure that an adequate dosage is administered to the patient.

One approach to improving the anti-settling properties of a suspension is to use a viscosity agent such as any of the natural gums or cellulose, such as hydroxypropylcellulose (HPC) or hydroxypropylmethylcellulose (HPMC) to increase viscosity, and thereby retard the rate of re-settling of wetted particles in the suspension. Such an approach has been found to be problematic in the case of ziprasidone hydrochloride since, once a viscosity agent has been added, when the ziprasidone hydrochloride eventually settles, it tends to form a thin cake which sits on the bottom and can be very difficult to break up and to re-suspend. Such caking is facilitated by temperature fluctuations and by vibrations such as those which occur during normal handling and transportation.

Further, in the specific case of ziprasidone acid addition salts, such salts generally exhibit a very bitter taste, the degree of bitterness increasing with increasing solubility of the particular salt. Sugars, with or without the presence of other sweetening and/or flavoring agents, are generally insufficient to mask the bitter

taste. Adjusting the pH to form the less soluble, hence less bitter, free base, is an option for decreasing bitterness. However, such adjustment can lead to changes in particle size if very careful and continuous control is not maintained. Substantial changes in particle size can, in turn, undesirably lead to changes in bioavailability.

5 Thus, a suspension comprising ziprasidone free base or ziprasidone hydrochloride (or other pharmaceutically acceptable ziprasidone addition salt) which maintains an improved shelf life (i.e., which maintains a longer period of suspension prior to re-settling) and which is easily re-suspendible would represent a valuable addition to the formulations arts. In the particular case of a suspension of a
10 ziprasidone acid addition salt, a suspension with improved taste would be a further valuable addition.

Summary Of The Invention

15 It has now been determined that dry components comprising ziprasidone free base or a difficult to wet pharmaceutically acceptable ziprasidone acid addition salt, a polysorbate, a viscosity agent, and colloidal silica easily form a good suspension in aqueous media. Thus one aspect of the invention is a composition of matter which is a suspension comprising ziprasidone free base or a difficult to wet pharmaceutically acceptable ziprasidone acid addition salt, water, a polysorbate, a viscosity agent,
20 and colloidal silicon dioxide. By "difficult to wet" it is meant a pharmaceutically acceptable salt of ziprasidone that does not readily form a suspension in water when mixed by ordinary means such as a laboratory blender which has been run at normal speed for ten minutes. Such salts do not, by themselves, form a "good suspension", as defined below, in water.

25 Again, it is noted ziprasidone hydrochloride is used herein to exemplify the invention, although it is to be understood that the use of an example is not to be considered as limiting.

The particle size of the ziprasidone hydrochloride particles is not considered particularly important in terms of ability to be wetted, although the mean particle size
30 is generally below or equal to 85 μm .

Pre-constituted suspensions according to the invention exhibit good suspension and can be remixed upon settling. By "good suspension" it is meant (1) that in a suspension according to the invention there is no visible settling for greater than 24 hours at room temperature (RT, usually 25°C), preferably for greater than

one week and (2) that when visible settling does occur, resuspension is easily effected by simple physical mixing such as gentle manual stirring or moderate manual shaking, high shear mixing not being required.

5 The invention is surprising in that polysorbates are the only wetting agents among a range of wetting agents that were tested which yielded positive results. Other wetting agents such as sodium lauryl sulfate caused foaming before adequate wetting could be obtained. Such foaming was problematic in that, although some particles appeared to wet, other salt particles became entrained in the foam, and still other salt particles sat as a dry mass atop the water, with the result that a
10 homogeneous suspension did not form. Thus, polysorbates are useful as agents in the instant invention that allowed complete wetting below the level at which they foam.

The term "polysorbate" is employed for its art-recognized meaning, i.e., polyoxyethylene sorbitan fatty acid esters as disclosed and defined in the Handbook
15 Of Pharmaceutical Excipients, edited by Ainley Wade and Paul Weller, The Pharmaceutical Press, London, 1994. Useful polysorbates include polysorbate 20, 21, 40, 60, 61, 65, 80, 81, 85, and 120. Polysorbate 80 is preferred.

The colloidal silicon dioxide useful herein is of the type known in the art (e.g., of the type available commercially as CAB-O-SIL[®], registered trademark of Cabot
20 Corporation, Boston, MA) and, while not wishing to be bound by any particular theory or mechanism, is believed to function as an anti-caking agent. That is, even though some re-settling may occur in a suspension according to the invention, the re-settled material does not cake, meaning that, even though a viscosity agent is present, resettled material does not form a firm mass that is difficult to break up and
25 re-disperse in the presence of colloidal silicon dioxide. In the presence of colloidal silicon dioxide, anti-caking properties are effected such that re-suspension can be effected by simple physical mixing, as previously described. Thus the advantage of the invention is that the combination of a polysorbate and colloidal silicon dioxide achieves good wetting together with facile re-suspendability in the event settling
30 occurs.

In a more specific aspect of the invention, there is provided an aqueous suspension of ziprasidone comprising a ziprasidone acid addition salt suspended in an aqueous pharmaceutically acceptable carrier which contains an alkali metal (e.g., sodium, lithium, or potassium) chloride or an alkaline earth metal (e.g. magnesium

or calcium) chloride. It has been determined that these salts are very effective as taste masking agents. While not wishing to be bound by theory or statements of mechanism, it is believed that such salts reduce the solubility, and thereby the dissolved amount and/or concentration, of ziprasidone salt in the suspension.

- 5 Reduced concentration of dissolved ziprasidone is not a disadvantage in a suspension dosage form, however, and in the instant invention is advantageous since suspended ziprasidone is more chemically stable than dissolved ziprasidone, thereby improving chemical stability, as well as taste.

10 The invention further provides a method of treating a psychosis, comprising administering to a patient in need of such treatment an effective amount of a suspension comprising ziprasidone free base or a pharmaceutically acceptable ziprasidone acid addition salt, a polysorbate, a viscosity agent, colloidal silicon dioxide, and water, said suspension being otherwise disclosed and described herein.

15 Ziprasidone hydrochloride can be used in any active crystalline or amorphous form, although crystalline ziprasidone hydrochloride monohydrate is preferred.

Detailed Description

20 In general, the polysorbate is used in an amount which is at least sufficient to effect complete wetting, i.e., meaning that drug is easily dispersed within a reasonable amount of time. The amount of polysorbate employed should not, as an upper limit, equal or exceed the amount which causes foaming.

25 Colloidal silicon dioxide is used in at least an anti-caking amount, i.e., an amount such that a firm, difficult to re-suspend cake does not form under normal conditions of transportation and storage. It is possible that re-settling may occur which is easily re-suspended by mild physical agitation (e.g., stirring or shaking). The upper limit for colloidal silicon dioxide is an amount above which, in conjunction with the viscosity agent, causes gelation. The particular grade of colloidal silicon dioxide, so long as it is pharmaceutically acceptable, has not been found to be critical.

30 Both the amount of polysorbate and colloidal silicon dioxide are system specific, optimum amounts depending on the drug form (e.g., free base versus acid salt), and the amounts and types of other excipients which may also be used as part of the suspension. Generally, the polysorbate will be employed in an amount of from about 0.01 to about 2.0 by weight % based on the weight of the final suspension,

more preferably from about 0.05 to about 0.30 weight %. The colloidal silicon dioxide will generally be employed in an amount of from 0.05 weight % to 2.0 weight % based on the weight of the suspension.

A suitable viscosity agent (also referred to in the art as a "thickening agent") is also used as a component of the invention. Such viscosity agents function as suspending agents and include, for example, hydrocolloid gums known for such purpose, examples of which include xanthan gum, guar gum, locust bean gum, gum tragacanth, and the like. Alternatively, synthetic suspending agents may be used such as sodium carboxymethylcellulose, polyvinylpyrrolidone, hydroxypropylcellulose, hydroxypropylmethylcellulose, and the like. The viscosity agent is generally used in an amount of from about 0.01 weight % to about 10 weight % based on the weight of the suspension. The amount actually used in a particular formulation is dependent on the exact agent and on other excipients present.

Ziprasidone free base or an acid addition salt can be used in this invention in any form, including anhydrous or hydrated. The ziprasidone hydrochloride employed herein, including the examples, was ziprasidone hydrochloride monohydrate, and is generally referred to throughout simply as ziprasidone hydrochloride for convenience. The invention is applicable to other ziprasidone acid addition salts as well, such as acetic, lactic, succinic, maleic, tartaric, citric, gluconic, ascorbic, benzoic, cinnamic, fumaric, sulfuric, phosphoric, hydrobromic, hydroiodic, sulfamic, sulfonic such as methanesulfonic, benzenesulfonic, and related acids.

As previously stated, it is advantageous to include a taste masking agent in the suspension in the specific case of pharmaceutically acceptable ziprasidone acid addition salts. Such taste masking agents are alkali metal and alkaline earth metal chlorides including sodium chloride, lithium chloride, potassium chloride, magnesium chloride, and calcium chloride. Sodium chloride is preferred. The taste masking agent is generally included in the suspension in a taste-masking amount, generally an amount of about 0.5 to about 2.0 weight % as sodium chloride based on the weight of the suspension. For other salts, equivalent molar amounts can be calculated.

A composition according to the invention is an oral, pre-constituted suspension which contains, as necessary ingredients, ziprasidone free base or a ziprasidone acid addition salt, water, a polysorbate, a viscosity agent, and colloidal silicon dioxide. Compositions according to the invention can also contain other

conventional pharmaceutically acceptable excipients such as, for example: flavorings, buffers, pH adjusting agents, diluents, colors, preservatives, and sweetening agents. Some excipients can serve multiple functions.

Flavors incorporated in the composition may be chosen from synthetic flavor oils and flavoring aromatics and/or natural oils, extracts from plant leaves, flowers, fruits, and so forth and combinations thereof. These may include cinnamon oil, oil of wintergreen, peppermint oils, clove oil, bay oil, anise oil, eucalyptus, thyme oil, cedar leaf oil, oil of nutmeg, oil of sage, oil of bitter almonds, and cassia oil. Also useful as flavors are vanilla, citrus oil, including lemon, orange, grape, lime and grapefruit, and fruit essences, including apple, banana, pear, peach, strawberry, raspberry, cherry, plum, pineapple, apricot, and so forth. The amount of flavoring may depend on a number of factors including the organoleptic effect desired. Generally the flavoring will be present in an amount of from about 0.01 to about 1.0 percent by weight based on the total suspension weight, when a flavor is used.

The suspensions can be manufactured by conventional pharmaceutical methods, combining the various components using conventional equipment such as an overhead stirrer, usually at about 100-500 RPM. Many different orders of adding components to the stirrer can be employed. A typical order of mixing in the instant invention, although others are certainly possible, is (1) adding water heated to 70°C for any components which require temperatures higher than room temperature (rt) as an aid to dissolution (if no such ingredients are employed, then heating is not required); (2) cooling to rt (about 30°C); then adding, in the following order (and assuming each component is employed), viscosity agent, sweetener, buffer, polysorbate, taste masking agent if employed, ziprasidone, colloidal silicon dioxide, and flavors.

The invention is further exemplified and disclosed by the following non-limiting examples:

EXAMPLE 1

A suspension formulation was prepared by heating 733.31 g of water to 70 °C followed by adding 1.36 g methylparaben and 0.17 g propylparaben while stirring at about 200 rpm with an overhead stirrer. After the parabens completely dissolved, the temperature was lowered to about 30°C. The following components were then added in order: 2.78 g xanthan gum, 333.90 g xylitol, 1.13 g anhydrous citric acid, 1.21 g trisodium citrate dihydrate, 0.55 g polysorbate 80, 11.13 g NaCl, 11.33 g ziprasidone hydrochloride monohydrate having a nominal particle size of 38 µm, 11.13 g colloidal silicon dioxide, and 5.0 g cherry flavor. The pH was adjusted to 4.0 using aqueous sodium hydroxide and hydrochloric acid as needed.

Example 2

This example discloses a process for making a ziprasidone free base suspension.

Into a 2 liter beaker was weighed 812.9 g of water which was stirred using an overhead stirrer at a speed of about 200 rpm. The water was heated to 70°C. Once the temperature reached 70°C, 1.36 g of methylparaben and 0.17 g of propylparaben were added. When the parabens were completely dissolved, the temperature was lowered to 40°C. To the solution was slowly added 3.27 g of a viscosity agent, CARBOPOL[®] resin 974P (Union Carbide Corporation, Danbury, CT), taking care to avoid big lumps, and increasing the stirring speed as necessary. Agitation was maintained until the viscosity agent had completely dispersed and/or dissolved. To the solution was added 218 g of sucrose. After dissolving the sucrose, temperature was lowered to 30°C. To the solution was added 2.94 g of trisodium citric salt. To the solution was added 0.544 g of polysorbate 80. To the solution was slowly added 11.325 g of ziprasidone free base. A 10% NaOH solution was used to adjust the pH of the formulation to 5.7. After the pH had equilibrated, 1.09 g of colloidal silicon dioxide (CAB-O-SIL[®], Cabot Corporation) was added.

EXAMPLE 3

This example illustrates the results obtained using colloidal silicon dioxide as an anti-caking agent as compared to other agents used for the same purpose.

Ziprasidone free base suspensions (nominal ziprasidone particle size of 38 μm) containing the same components, except for the anti-caking agent, were made as in Example 2, with and without each of the anti-caking agents listed below. Each anti-caking agent was added into the formulation in a 60 cc bottle, which was then centrifuged at 2000 RPM for 20 minutes to accelerate settling, and the bottles were then gently rotated at a speed of about one rotation every 2 seconds to resuspend the settled solids. The time required for the formulations to become fully homogeneous (no solids stuck at the bottom of the bottle by visual inspection) was recorded. The data support that the time required for the formulations to resuspend is substantially reduced using colloidal silicon dioxide (CSD in the Table below), reducing resuspension time substantially at about 0.3% level.

No.	Anti-caking Agent	Level (w/w %) in Formulation	Resuspendability
1	None		> 20 minutes
2	MCC, 200 μm	3.0	> 20 minutes
3	MCC, 50 μm	1.0	> 20 minutes
4	SiO ₂	1.0	~ 2 minutes
5	CSD	0.3	~ 1 minutes
6	Magnesium Stearate	0.2	> 20 minutes

MCC is an acronym for microcrystalline cellulose

Particle sizes, given in microns, are nominal

Example 4

Different surfactants were tested for their ability to wet ziprasidone HCl monohydrate in water: sodium lauryl sulfate (SLS), MIGLYOL[®] (registered trademark of Dynamit Nobel Aktiengesellschaft, Germany) Triglyceride (810), and polysorbate-80. In three separate 100 mL volumes of water, each surfactant listed above was added in an amount to make a 1% solution. MIGLYOL[®] was added in excess due to its low solubility. While stirring at 200 rpm with an overhead stirrer, to each of the three surfactant solutions was added 1.132 g of ziprasidone was added to each of the three surfactant solutions (equivalent to 10 mgA/ml). The observed rate of wetting was recorded as follows:

- | | |
|--------------------------------------|----------|
| 1) SLS | > 20 min |
| 2) MIGLYOL [®] triglyceride | >20 min |
| 3) Polysorbate-80 | < 2 min |

Only the polysorbate 80 solution did not show any visible aggregates after mixing overnight.

Further testing of polysorbate-80 at different concentrations demonstrated that a concentration as low as 0.05% can significantly decrease drug wetting time.

<u>Conc. (%)</u>	<u>Drug wetting time</u>
0.05	4 min
0.15	3.8 min
0.25	3.5 min

What is claimed is:

1. A composition comprising ziprasidone free base or a pharmaceutically acceptable ziprasidone acid addition salt, water, a polysorbate, a viscosity agent, and colloidal silicon dioxide.
- 5 2. A composition as defined in claim 1, wherein said ziprasidone acid addition salt is ziprasidone hydrochloride.
3. A composition as defined in claim 1, wherein said polysorbate is polysorbate 20, 21, 40, 60, 61, 65, 80, 81, 85, or 120.
4. A composition as defined in claim 3, wherein said polysorbate is
10 polysorbate 80.
5. A composition as defined in claim 1, further comprising a taste masking agent selected from alkali metal chlorides and alkaline earth metal chlorides.
6. A composition as defined in claim 5, wherein said alkali metal salt is sodium chloride, potassium chloride, or lithium chloride.
- 15 7. A composition as defined in claim 6, wherein said alkali metal salt is sodium chloride.
8. A composition as defined in claim 5, wherein said alkaline earth metal salt is magnesium chloride or calcium chloride.
9. A composition comprising ziprasidone free base or ziprasidone
20 hydrochloride, a polysorbate, colloidal silicon dioxide, a viscosity agent, and water.
10. A composition as defined in claim 9, wherein said polysorbate is polysorbate 20, 21, 40, 60, 61, 65, 80, 81, 85, or 120.
11. A composition as defined in claim 10, wherein said polysorbate is polysorbate 80.
- 25 12. A composition as defined in claim 9, further comprising a taste masking agent selected from alkali metal chlorides and alkaline earth metal chlorides.
13. A composition as defined in claim 12, wherein said alkali metal salt is sodium chloride, potassium chloride, or lithium chloride.
14. A composition as defined in claim 13, wherein said alkali metal salt is
30 sodium chloride.
15. A composition as defined in claim 12, wherein said alkaline earth metal salt is magnesium chloride or calcium chloride.

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/IB 00/00593

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/496 A61K9/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X, L	EP 0 965 343 A (PFIZER PRODUCTS INC.) 22 December 1999 (1999-12-22) page 9; examples 11,12 the priority claim of the present application might not be justified	1-14

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

17 August 2000

Date of mailing of the international search report

24/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Benz, K

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 00/00593

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 965343 A	22-12-1999	AU 3398399 A	23-12-1999
		HR 990193 A	29-02-2000
		JP 2000007566 A	11-01-2000
		NO 992892 A	16-12-1999
		PL 333737 A	20-12-1999